

Spectrum of boundary states in N=1 SUSY sine-Gordon theory

Z. Bajnok*, L. Palla† and G. Takács‡

Institute for Theoretical Physics
Eötvös University
H-1117 Budapest, Pázmány Péter sétány 1/A

25 June 2002

Abstract

We consider N=1 supersymmetric sine-Gordon theory (*SSG*) with supersymmetric integrable boundary conditions (boundary *SSG=BSSG*). We find two possible ways to close the boundary bootstrap for this model, corresponding to two different choices for the boundary supercharge. We argue that these two bootstrap solutions should correspond to the two integrable Lagrangian boundary theories considered recently by Nepomechie.

PACS numbers: 11.55.Ds, 11.30.Pb, 11.10.Kk

Keywords: supersymmetry, sine-Gordon model, integrable quantum field theory, boundary scattering, bootstrap

1 Introduction

In this paper we consider the N=1 supersymmetric sine-Gordon theory with supersymmetric integrable boundary conditions (*BSSG*). Our aim is to find a closure of the boundary bootstrap for this model.

*bajnok@afavant.elte.hu

†palla@ludens.elte.hu

‡takacs@ludens.elte.hu

The $N=1$ supersymmetric sine-Gordon model is the natural supersymmetric extension of the ordinary sine-Gordon model. It is an integrable field theory with infinitely many conserved charges [1]. The S matrix of the theory was obtained in [2], while the integrable and supersymmetric boundary conditions were considered in [3], but it took a while until the most general integrable supersymmetric boundary interaction was found [4].

As a result of integrability, the boundary scattering factorizes, and the general solution of the boundary Yang-Baxter equation was found in [5], but the constraint of supersymmetry was not imposed. Nepomechie was the first to consider supersymmetric boundary scattering [4], building on previous results obtained in the case of supersymmetric sinh-Gordon theory [6]. However, no one has exposed the full structure of the solitonic reflection amplitude, although the results obtained in the case of the tricritical Ising model [7] are closely related to this problem. Besides that, the closure of the bootstrap and the spectrum of boundary states have not been even touched before. Therefore our aim is to clear up the issue of supersymmetric boundary scattering in the $BSSG$ model and to find the complete spectrum of boundary states and their associated reflection factors. The main idea – motivated by the successful description of the bulk scattering – is to look for the reflection amplitudes in a form where there is no mixing between the supersymmetric and other internal quantum numbers. This means an Ansatz for the reflection amplitudes as a product of two terms one of which is the ordinary (bosonic) sine-Gordon reflection amplitude, while the other describes the scattering of the SUSY degrees of freedom.

Within the bootstrap procedure, we consider first the solitonic reflection amplitudes on the ground state and on the first two excited boundaries. The SUSY factors in these solutions have no poles in the physical strip, thus the masses of boundary states emerging are the same as in the bosonic theory, however, SUSY introduces a nontrivial degeneracy. The spectrum of general higher excited boundaries is easily extracted from these results. We determine also the various reflection amplitudes on these excited boundaries.

There are two ways to close the bootstrap, starting from two different ground state reflection amplitudes, corresponding to two possible choices of the boundary supercharge. Both solutions lead to the same spectrum of boundary states, but the reflection amplitudes are different. In one case the reflections conserve fermionic parity, while in the other they do not.

The layout of the paper is as follows. Section 2 recalls briefly some important facts about supersymmetric sine-Gordon theory. In Section 3 the reader is reminded of the supersymmetric and integrable boundary interactions that can be added to the theory, and we also discuss of the boundary supercharge and derive a formula relating it to the boundary Hamiltonian. Section 4 gives a quick review (containing only the most necessary facts) of the spectrum and reflection factors of the ordinary (non supersymmetric) sine-Gordon model with integrable boundary conditions. In Section 5 we present the main results of the paper, which is a conjecture for the spectrum and the full set of reflection factors of the $BSSG$ model. We consider the breather reflection amplitudes in Section 6 and then give our conclusions in Section 7.

2 Bulk SSG theory

The bulk supersymmetric sine-Gordon theory is defined by the classical action

$$\begin{aligned}\mathcal{A}_{SSG} &= \int dt dx \mathcal{L}_{SSG}(x, t) \\ \mathcal{L}_{SSG} &= \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi + i \bar{\Psi} \gamma^\mu \partial_\mu \Psi + m \bar{\Psi} \Psi \cos \frac{\beta}{2} \Phi + \frac{m^2}{\beta^2} \cos \beta \Phi\end{aligned}\quad (1)$$

where Φ is a real scalar, Ψ is a Majorana fermion field, m is a mass parameter and β is the coupling constant. The theory is invariant under an $N = 1$ supersymmetry algebra and has infinitely many commuting local conserved charges [1]. These charges survive at the quantum level and render the theory integrable, which makes it possible to describe the exact spectrum and the S matrix.

2.1 Spectrum and bulk scattering amplitudes

The spectrum consists of the soliton/antisoliton multiplet, realizing supersymmetry in a nonlocal way, and breathers that are bound states of a soliton with an antisoliton.

The building blocks of supersymmetric factorized scattering theory were first described in [10], using an Ansatz in which the full scattering amplitude is a direct product of a part carrying the SUSY structures and a part describing all the rest of the dynamics. The full SSG S matrix was constructed in [2].

The supersymmetric solitons are described by RSOS kinks $K_{ab}^\epsilon(\theta)$ of mass M and rapidity θ , where a, b take the values $0, \frac{1}{2}$ and 1 with $|a - b| = 1/2$, and describe the supersymmetric structure, while $\epsilon = \pm$ corresponds to topological charge ± 1 (soliton/antisoliton). Multi-particle asymptotic states are built as follows

$$\left| K_{a_0 a_1}^{\epsilon_1}(\theta_1) K_{a_1 a_2}^{\epsilon_2}(\theta_2) \dots K_{a_{N-1} a_N}^{\epsilon_N}(\theta_N) \right\rangle \quad (2)$$

where $\theta_1 > \theta_2 > \dots > \theta_N$ for an *in* state and $\theta_1 < \theta_2 < \dots < \theta_N$ for an *out* state. The two-particle scattering process

$$K_{ab}^{\epsilon_1}(\theta_1) + K_{bc}^{\epsilon_2}(\theta_2) \rightarrow K_{ad}^{\epsilon'_2}(\theta_2) + K_{dc}^{\epsilon'_1}(\theta_1)$$

has an amplitude of the form

$$S_{SUSY} \left(\begin{array}{cc} a & d \\ b & c \end{array} \middle| \theta_1 - \theta_2 \right) \times S_{SG}(\theta_1 - \theta_2)_{\epsilon_1 \epsilon_2}^{\epsilon'_1 \epsilon'_2} \quad (3)$$

i.e. the tensor structure of the scattering amplitude factorizes into a part describing the SUSY structure (which we call the SUSY factor) and another part corresponding to the topological charge (the bosonic factor).

The bosonic factor coincides with the usual sine-Gordon S matrix¹

$$\begin{aligned}
S_{SG}(u)_{++}^{++} &= S_{SG}(u)_{--}^{--} = - \prod_{l=1}^{\infty} \left[\frac{\Gamma(2(l-1)\lambda - \frac{\lambda u}{\pi}) \Gamma(2l\lambda + 1 - \frac{\lambda u}{\pi})}{\Gamma((2l-1)\lambda - \frac{\lambda u}{\pi}) \Gamma((2l-1)\lambda + 1 - \frac{\lambda u}{\pi})} / (u \rightarrow -u) \right] \\
S_{SG}(u)_{+-}^{+-} &= S_{SG}(u)_{-+}^{-+} = \frac{\sin(\lambda u)}{\sin(\lambda(\pi - u))} S_{SG}(u)_{++}^{++} \quad ; \quad \lambda = \frac{2\pi}{\beta^2} - \frac{1}{2} \\
S_{SG}(u)_{-+}^{-+} &= S_{SG}(u)_{+-}^{+-} = \frac{\sin(\lambda\pi)}{\sin(\lambda(\pi - u))} S_{SG}(u)_{++}^{++} \quad ; \quad u = -i\theta \quad , \quad (4)
\end{aligned}$$

while the SUSY factor is identical to the S matrix of the tricritical Ising model perturbed by the primary field of dimension $\frac{3}{5}$ [8]:

$$\begin{aligned}
S_{SUSY} \left(\begin{array}{c|c} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{array} \middle| \theta \right) &= S_{SUSY} \left(\begin{array}{c|c} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{array} \middle| \theta \right) = 2^{(i\pi-\theta)/2\pi i} \cos\left(\frac{\theta}{4i} - \frac{\pi}{4}\right) K(\theta) \\
S_{SUSY} \left(\begin{array}{c|c} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{array} \middle| \theta \right) &= S_{SUSY} \left(\begin{array}{c|c} \frac{1}{2} & 1 \\ 1 & \frac{1}{2} \end{array} \middle| \theta \right) = 2^{\theta/2\pi i} \cos\left(\frac{\theta}{4i}\right) K(\theta) \\
S_{SUSY} \left(\begin{array}{c|c} 0 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{array} \middle| \theta \right) &= S_{SUSY} \left(\begin{array}{c|c} 1 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{array} \middle| \theta \right) = 2^{(i\pi-\theta)/2\pi i} \cos\left(\frac{\theta}{4i} + \frac{\pi}{4}\right) K(\theta) \\
S_{SUSY} \left(\begin{array}{c|c} \frac{1}{2} & 1 \\ 0 & \frac{1}{2} \end{array} \middle| \theta \right) &= S_{SUSY} \left(\begin{array}{c|c} \frac{1}{2} & 0 \\ 1 & \frac{1}{2} \end{array} \middle| \theta \right) = 2^{\theta/2\pi i} \cos\left(\frac{\theta}{4i} - \frac{\pi}{2}\right) K(\theta)
\end{aligned}$$

$$K(\theta) = \frac{1}{\sqrt{\pi}} \prod_{k=1}^{\infty} \frac{\Gamma(k - 1/2 + \theta/2\pi i) \Gamma(k - \theta/2\pi i)}{\Gamma(k + 1/2 - \theta/2\pi i) \Gamma(k + \theta/2\pi i)}$$

As the SUSY factor has no poles in the physical strip, the solitonic amplitudes (3) have poles at exactly the same locations as the sine-Gordon soliton S matrix. These correspond to bound states (breathers) $\mathbf{B}_n$ of mass

$$m_n = 2M \sin \frac{\pi n}{2\lambda} \quad , \quad n = 1, \dots [\lambda] \quad .$$

The S matrix of the breathers was first found in [9]. The breathers form a particle multiplet composed of a boson and a fermion, on which supersymmetry is represented in a standard way [10, 11].

For the ordinary sine-Gordon theory, the correspondence between the Lagrangian theory and the bootstrap S matrix (4) is very well established. There is much less evidence for the correctness of the S matrix (3) as the scattering amplitude of SUSY sine-Gordon theory. Besides the original construction [2] (based on arguments related to $N = 1$ supersymmetric minimal models), another indication is that at a particular value of the coupling β where it is expected to have a restriction to the SUSY version of Lee-Yang theory (superconformal minimal model $\mathcal{SM}(2/8)$ perturbed by the relevant superconformal primary

¹Note that the relation between the parameter λ and the coupling β is different from the sine-Gordon case.

field $\Phi(1, 3)$, which is equivalent to Virasoro minimal model $\mathcal{M}(3/8)$ perturbed by the primary field $\Phi(1, 5)$, the first breather supermultiplet has the same scattering amplitude as predicted from RSOS restriction of imaginary coupled $a_2^{(2)}$ Toda theory in [12] (see also [13]).

2.2 Bulk SUSY charges

The bulk theory has two supersymmetry charges of opposite chirality Q and $\bar{Q}$, which together form a Majorana spinor. They act on one-particle states $|A_i(\theta)\rangle$ in the following way [8, 10, 11]:

$$Q|A_i(\theta)\rangle = \sqrt{m_i}e^{\theta/2}\mathcal{Q}|A_i(\theta)\rangle \quad , \quad \bar{Q}|A_i(\theta)\rangle = \sqrt{m_i}e^{-\theta/2}\bar{\mathcal{Q}}|A_i(\theta)\rangle$$

where m_i are the particle masses and $\mathcal{Q}$, $\bar{\mathcal{Q}}$ are matrices satisfying

$$\mathcal{Q}^2 = 1 \quad , \quad \bar{\mathcal{Q}}^2 = 1 \quad .$$

In the one-particle basis $\{K_{0\frac{1}{2}}, K_{1\frac{1}{2}}, K_{\frac{1}{2}0}, K_{\frac{1}{2}1}\}$ (we omit the upper index ϵ , as the SUSY action does not depend on the topological charge), the supersymmetry algebra is represented by the matrices

$$\mathcal{Q} = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad , \quad \bar{\mathcal{Q}} = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

The SUSY algebra of the sine-Gordon theory has a central charge and the matrix

$$Z = \frac{1}{2}\{\mathcal{Q}, \bar{\mathcal{Q}}\} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

describes the SUSY central charge in the above basis. This is not to be confused with the topological charge T of the sine-Gordon solitons, which is represented by the upper indices ϵ . Z can take the values 0 or ± 1 , and it distinguishes between solitons/antisolitons mediating from odd to even and from even to odd vacua of the bosonic potential [14] (this means that using the terminology of [15] the theory is 2-folded).

The above representation of SUSY describes BPS saturated objects. There was a controversy in the literature whether the solitons in SSG are BPS saturated, since $N = 1$ SUSY does not protect their mass from acquiring radiative corrections. However, it was shown in [16] that they remain BPS saturated at one-loop and probably to all orders, due to anomalous quantum corrections to the classical formula for the central charge Z .

The fermionic parity operator $\Gamma = (-1)^F$ is given by the matrix

$$\Gamma = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

Using the definition of Γ it is possible to specify a basis of pure bosonic and pure fermionic states for any given (fixed) number of particles. However, the composition (coproduct) rules of the kink states as given in (2) are not free and therefore in the boson-fermion internal space supersymmetry acts nonlocally. The action of supersymmetry on multi-particle states involves braiding factors depending on Γ that are defined by the coproduct Δ :

$$\begin{aligned} \Delta(Q) &= Q \otimes \mathbb{I} + \Gamma \otimes Q \\ \Delta(\bar{Q}) &= \bar{Q} \otimes \mathbb{I} + \Gamma \otimes \bar{Q} \\ \Delta(\Gamma) &= \Gamma \otimes \Gamma \end{aligned}$$

The action on breather states can be derived using the bootstrap, but can also be obtained from the representation theory of the SUSY algebra. It turns out that the central charge Z (as well as the topological charge T) vanishes identically for the breathers. For further details we refer to [10, 11].

3 Boundary SSG theory

3.1 Integrable and supersymmetric boundary interactions

Boundary *SSG* (*BSSG*) theory can be described by an action of the form

$$\mathcal{A}_{BSSG} = \int_{-\infty}^{\infty} dt \int_{-\infty}^0 dx \mathcal{L}_{SSG}(x, t) + \int_{-\infty}^{\infty} L_B(t) dt$$

where $L_B(t)$ is a term local at $x = 0$. We consider boundary interactions which are both supersymmetric and integrable. The case when $L_B(t)$ depends only on the value of Φ and Ψ at $x = 0$ was considered in [3], where it was found that supersymmetry and integrability restricts the boundary interaction to the form

$$L_B = \pm \frac{4m}{\beta^2} \cos \frac{\beta}{2} \Phi \pm \bar{\psi} \psi \Big|_{x=0} \quad (5)$$

which gives four discrete choices. Here we write the Majorana spinor in component form

$$\Psi = \begin{pmatrix} \bar{\psi} \\ \psi \end{pmatrix}.$$

However, this is not the whole story. The Majorana fermion in the ultraviolet limit is described by the $c = \frac{1}{2}$ Ising conformal field theory. It was noted in [17] that in order for the Majorana fermion to describe correctly the boundary states of the conformal field theory, one has to include a fermionic boundary degree of freedom $a(t)$ that is related to the boundary value of the Ising spin. Using this fact, a two-parameter set of integrable supersymmetric boundary conditions was derived in [4]. The boundary interaction term is of the form

$$L_B^\pm = (\pm \bar{\psi} \psi + ia \partial_t a - 2f(\Phi) a(\psi \mp \bar{\psi}) + \mathcal{B}(\Phi))|_{x=0} \quad (6)$$

The functions f and $\mathcal{B}$ are fixed by the requirement of boundary integrability and supersymmetry (cf. [4]). We denote the theories obtained by adding $L_B^\pm$ to the bulk action as $BSSG^\pm$. The most important property of this action is that it depends on two continuously varying boundary parameters, exactly as in the case of the non-supersymmetric boundary sine-Gordon theory [18]. This is important for consistency with the bootstrap since the reflection factors we find depend on two parameters as well. The boundary Lagrangian (5) can be obtained as a special case of (6) when the parameters are tuned so that $f = 0$, and therefore the boundary fermion a decouples.

3.2 The boundary SUSY charge and the Hamiltonian

The bulk SUSY charges Q and $\bar{Q}$ can be written as integrals of local (fermionic) densities q and $\bar{q}$:

$$Q = \int_{-\infty}^{\infty} q(x, t) dx \quad , \quad \bar{Q} = \int_{-\infty}^{\infty} \bar{q}(x, t) dx$$

They have the anticommutation relation

$$\{Q, \bar{Q}\} = 2M Z$$

and satisfy the following relation (among others)

$$QQ + \bar{Q}\bar{Q} = 2H \quad (7)$$

where $H = \int h(x, t) dx$ is the Hamiltonian, Z is the SUSY central charge and M is the soliton mass. In a boundary theory with supersymmetric integrable boundary condition, the conserved supercharge can be written as follows:

$$\tilde{Q}_\pm = \int_{-\infty}^0 (q(x, t) \pm \bar{q}(x, t)) dx + Q_B(x = 0, t)$$

where Q_B is a boundary contribution, localized at $x = 0$. There are two possible choices ($\pm$) corresponding to the two possible conformal fermionic boundary conditions in the ultraviolet limit. As shown in [4], these correspond to the two choices of sign in (6) and therefore to $BSSG^\pm$.

Similarly, the Hamiltonian takes the form

$$\tilde{H} = \int_{-\infty}^0 h(x, t) dx + H_B(x = 0, t)$$

where H_B is the boundary interaction. Let us for the moment neglect the contribution from the central charge Z (this is possible e.g. in a sector containing only breathers). Then, using eqn. (7) it is easy to see that

$$\tilde{Q}_{\pm}^2 = 2 \int_{-\infty}^0 h(x, t) dx + 2H'_B(x = 0, t)$$

where H'_B is again some term localized at $x = 0$. However, $\tilde{Q}_{\pm}$ is conserved in time, and therefore

$$\frac{d}{dt} \left(2 \int_{-\infty}^0 h(x, t) dx + 2H'_B(x = 0, t) \right) = 0.$$

In the supersymmetric boundary sine-Gordon theory, this property uniquely determines H'_B as the boundary interaction H_B and therefore

$$\tilde{Q}_{\pm}^2 = 2\tilde{H}$$

Including Z , it is natural to expect that this relation extends as follows:

$$\tilde{Q}_{\pm}^2 = 2 \left(\tilde{H} \pm M \tilde{Z} \right) \quad (8)$$

where $\tilde{Z}$ is an appropriate extension of Z to the boundary situation. We shall see that the two bootstrap solutions we propose correctly reproduce this formula.

4 Boundary sine-Gordon model

4.1 Ground state reflection factors

The most general reflection factor – modulo CDD-type factors – of the soliton antisoliton multiplet $|s, \bar{s}\rangle$ on the ground state boundary, denoted by $|\rangle$, satisfying the boundary versions of the Yang-Baxter, unitarity and crossing equations was found by Ghoshal and Zamolodchikov [18]:

$$\begin{aligned} R_{SG}(\eta, \vartheta, u) &= \begin{pmatrix} P^+(\eta, \vartheta, u) & Q(\eta, \vartheta, u) \\ Q(\eta, \vartheta, u) & P^-(\eta, \vartheta, u) \end{pmatrix} \\ &= \begin{pmatrix} P_0^+(\eta, \vartheta, u) & Q_0(u) \\ Q_0(u) & P_0^-(\eta, \vartheta, u) \end{pmatrix} R_0(u) \frac{\sigma(\eta, u)}{\cos(\eta)} \frac{\sigma(i\vartheta, u)}{\cosh(\vartheta)} \quad , \\ P_0^{\pm}(\eta, \vartheta, u) &= \cos(\lambda u) \cos(\eta) \cosh(\vartheta) \mp \sin(\lambda u) \sin(\eta) \sinh(\vartheta) \quad , \\ Q_0(u) &= -\sin(\lambda u) \cos(\lambda u) \quad , \end{aligned} \quad (9)$$

where η and ϑ are the two real parameters characterizing the solution,

$$R_0(u) = \prod_{l=1}^{\infty} \left[\frac{\Gamma(4l\lambda - \frac{2\lambda u}{\pi}) \Gamma(4\lambda(l-1) + 1 - \frac{2\lambda u}{\pi})}{\Gamma((4l-3)\lambda - \frac{2\lambda u}{\pi}) \Gamma((4l-1)\lambda + 1 - \frac{2\lambda u}{\pi})} / (u \rightarrow -u) \right]$$

is the boundary condition independent part and

$$\sigma(x, u) = \frac{\cos x}{\cos(x + \lambda u)} \prod_{l=1}^{\infty} \left[\frac{\Gamma(\frac{1}{2} + \frac{x}{\pi} + (2l-1)\lambda - \frac{\lambda u}{\pi}) \Gamma(\frac{1}{2} - \frac{x}{\pi} + (2l-1)\lambda - \frac{\lambda u}{\pi})}{\Gamma(\frac{1}{2} - \frac{x}{\pi} + (2l-2)\lambda - \frac{\lambda u}{\pi}) \Gamma(\frac{1}{2} + \frac{x}{\pi} + 2l\lambda - \frac{\lambda u}{\pi})} / (u \rightarrow -u) \right]$$

describes the boundary condition dependence. The reflection factors of the breathers can be obtained by the bulk bootstrap procedure [19].

4.2 The general spectrum and the associated reflection factors

The spectrum of boundary excited states was determined in [20, 21]. It can be parametrized by a sequence of integers $|n_1, n_2, \dots, n_k\rangle$, whenever the $\frac{\pi}{2} \geq \nu_{n_1} > w_{n_2} > \dots \geq 0$ condition holds, where

$$\nu_n = \frac{\eta}{\lambda} - \frac{\pi(2n+1)}{2\lambda} \quad \text{and} \quad w_k = \pi - \frac{\eta}{\lambda} - \frac{\pi(2k-1)}{2\lambda}.$$

The mass of such a state is

$$m_{|n_1, n_2, \dots, n_k\rangle} = M \sum_{i \text{ odd}} \cos(\nu_{n_i}) + M \sum_{i \text{ even}} \cos(w_{n_i}). \quad (10)$$

The reflection factors depend on which sectors we are considering. In the even sector, i.e. when k is even, we have

$$Q_{|n_1, n_2, \dots, n_k\rangle}(\eta, \vartheta, u) = Q(\eta, \vartheta, u) \prod_{i \text{ odd}} a_{n_i}(\eta, u) \prod_{i \text{ even}} a_{n_i}(\bar{\eta}, u),$$

and

$$P_{|n_1, n_2, \dots, n_k\rangle}^{\pm}(\eta, \vartheta, u) = P^{\pm}(\eta, \vartheta, u) \prod_{i \text{ odd}} a_{n_i}(\eta, u) \prod_{i \text{ even}} a_{n_i}(\bar{\eta}, u),$$

for the solitonic processes, where

$$a_n(\eta, u) = \prod_{l=1}^n \left\{ 2 \left(\frac{\eta}{\pi} - l \right) \right\} \quad ; \quad \bar{\eta} = \pi(\lambda + 1) - \eta$$

and

$$\{y\} = \frac{\left(\frac{y+1}{2\lambda}\right) \left(\frac{y-1}{2\lambda}\right)}{\left(\frac{y+1}{2\lambda} - 1\right) \left(\frac{y-1}{2\lambda} + 1\right)} \quad , \quad (x) = \frac{\sin\left(\frac{u}{2} + \frac{x\pi}{2}\right)}{\sin\left(\frac{u}{2} - \frac{x\pi}{2}\right)}$$

In the odd sector, i.e. when k is odd, the same formulae apply if in the ground state reflection factors the $\eta \leftrightarrow \bar{\eta}$ and $s \leftrightarrow \bar{s}$ changes are made. The breather sector can be obtained again by bulk fusion.

5 Supersymmetric boundary sine-Gordon model

5.1 Ground state reflection factors

5.1.1 The general solution for the reflection factor

Following the bulk case, we suppose that the reflection matrix factorizes as

$$R_{SUSY}(\theta) \times R_{SG}(\theta) .$$

In this special form the constraints as unitarity, boundary Yang-Baxter equation and crossing-unitarity relation [18] can be satisfied separately for the two factors. Since the sine-Gordon part already fulfills these requirements, we concentrate on the supersymmetric part.

From the RSOS nature of the bulk S -matrix (3) it is clear that the boundary must also have RSOS labels and the adjacency conditions between the nearest kink and the boundary must also hold. Thus the following reflections are possible:

$$K_{ba}(\theta)|B_a\rangle = \sum_c R_{ac}^b(\theta)K_{bc}(-\theta)|B_c\rangle$$

or in detail

$$K_{0\frac{1}{2}}(\theta)|B_{\frac{1}{2}}\rangle = R_{\frac{1}{2}\frac{1}{2}}^0(\theta)K_{0\frac{1}{2}}(-\theta)|B_{\frac{1}{2}}\rangle \quad ; \quad K_{1\frac{1}{2}}(\theta)|B_{\frac{1}{2}}\rangle = R_{\frac{1}{2}\frac{1}{2}}^1(\theta)K_{1\frac{1}{2}}(-\theta)|B_{\frac{1}{2}}\rangle .$$

and

$$K_{\frac{1}{2}a}(\theta)|B_a\rangle = R_{aa}^{\frac{1}{2}}(\theta)K_{\frac{1}{2}a}(-\theta)|B_a\rangle + R_{ab}^{\frac{1}{2}}(\theta)K_{\frac{1}{2}b}(-\theta)|B_b\rangle \quad , \quad b \neq a \quad , \quad a, b = 0, 1 \quad (11)$$

In the second process the label of the boundary state has changed, which shows that $|B_0\rangle$ and $|B_1\rangle$ form a doublet. All of the constraints mentioned above factorize in the sense that they give independent equations for the reflections on the boundary $|B_{1/2}\rangle$ and on the doublet $|B_{0,1}\rangle$. Since the ground state boundary is expected to be nondegenerate we first concentrate on reflection factors off the singlet boundary $|B_{1/2}\rangle$. The most general solution of the boundary Yang-Baxter equation is of the form [5]

$$R_{\frac{1}{2}\frac{1}{2}}^0(\theta) = (1 + A \sinh(\theta/2))M(\theta) \quad ; \quad R_{\frac{1}{2}\frac{1}{2}}^1(\theta) = (1 - A \sinh(\theta/2))M(\theta)$$

while unitarity and crossing symmetry give the following restrictions

$$M(\theta)M(-\theta)(1 - A^2 \sinh^2(\theta/2)) = 1 \quad ,$$

$$M\left(\frac{i\pi}{2} - \theta\right) = \cosh\left(\frac{\theta}{2} - \frac{i\pi}{4}\right) K(2\theta) 2^{(i\pi+2\theta)/2\pi i} M\left(\frac{i\pi}{2} + \theta\right) .$$

We suppose that the boundary states $|B_a\rangle$, $a = 0, 1$ can be obtained by boundary bootstrap from the ground state $|B_{1/2}\rangle$ [22, 7]. Therefore we do not consider the Yang-Baxter equation and the other constraints for the amplitudes $R_{ab}^{1/2}(\theta)$ as these will be guaranteed to be fulfilled by the bootstrap. We shall see later that in general the boundary states $|B_a\rangle$, $a = 0, 1$ come in multiple copies, each of which forms a doublet of states with the same energy.

5.1.2 Action of the boundary supersymmetry charges $\tilde{Q}_\pm$

We need to construct the action of the boundary supercharge on the asymptotic states. We expect that the action is given by

$$\tilde{Q}_\pm = Q \pm \bar{Q} + Q_B$$

where Q , $\bar{Q}$ act on the particles as in the bulk theory (Section 2.2)². The reason for this is that they are given by integrals of local (fermionic) densities, and asymptotic particles are localized far away from the wall, so the action of these charges is not affected by the presence of the boundary. Q_B is the action of the boundary contribution, which we take to be

$$Q_B = \gamma \Gamma, \quad (12)$$

where γ is some unknown parameter (related to the energy of the boundary ground state – see later). The reason for this choice is that we expect the boundary supercharge to commute with the bulk S -matrix, which is symmetric under the action of $Q, \bar{Q}$ and Γ by construction[10, 11]. (12) is also supported by classical considerations in [4], showing also that the classical version of γ is a function of the parameters in the boundary Lagrangian (6).

Next we need to give the action of $Q, \bar{Q}$ and Γ on the boundary ground state $|B_{1/2}\rangle$. Following [7], we choose

$$Q|B_{\frac{1}{2}}\rangle = 0 \quad , \quad \bar{Q}|B_{\frac{1}{2}}\rangle = 0 \quad , \quad \Gamma|B_{\frac{1}{2}}\rangle = |B_{\frac{1}{2}}\rangle. \quad (13)$$

The first two relations express that the boundary ground state is supersymmetric, while the last one shows that it is an eigenvector of Γ . We expect that because the ground state is nondegenerate. The choice of the eigenvalue (± 1) is not important, as it could be compensated by a redefinition of γ . It is a consequence of (8) and (13) that the ground state energy is $\gamma^2/2$, which will be shown later to be consistent with the action of $\tilde{Q}_\pm$ on the excited boundary states.

5.1.3 Supersymmetric reflection amplitudes

Now we would like to impose the supersymmetry constraints on the ground state reflection amplitudes.

The two choices $\tilde{Q}_\pm$ will give different solutions [7, 22]. If the boundary supercharge $\tilde{Q}_+$ commutes with the reflections (theory $BSSG^+$) then we obtain

$$R_{\frac{1}{2}\frac{1}{2}}^0(\theta) = R_{\frac{1}{2}\frac{1}{2}}^1(\theta) = 2^{-\theta/\pi i} \prod_{k=1}^{\infty} \left[\frac{\Gamma(k - \frac{\theta}{2\pi i})\Gamma(k - \frac{\theta}{2\pi i})}{\Gamma(k - \frac{1}{4} - \frac{\theta}{2\pi i})\Gamma(k + \frac{1}{4} - \frac{\theta}{2\pi i})} / \{\theta \leftrightarrow -\theta\} \right] = 2^{-\theta/\pi i} P(\theta) \quad (14)$$

²Note that our charge $\bar{Q}$ differs by a sign from that used in [7], while agrees with the convention in [11]. This is important when comparing our results with those in [7].

If, however, it is $\tilde{Q}_-$ that commutes with the reflections ($BSSG^-$) then the result is

$$\begin{aligned} R_{\frac{1}{2}\frac{1}{2}}^0(\theta) &= \left(\cos \frac{\xi}{2} + i \sinh \frac{\theta}{2} \right) K(\theta - i\xi) K(i\pi - \theta - i\xi) 2^{-\theta/\pi i} P(\theta) \\ R_{\frac{1}{2}\frac{1}{2}}^1(\theta) &= \left(\cos \frac{\xi}{2} - i \sinh \frac{\theta}{2} \right) K(\theta - i\xi) K(i\pi - \theta - i\xi) 2^{-\theta/\pi i} P(\theta) \end{aligned} \quad (15)$$

where ξ is related to γ as

$$\gamma = -2\sqrt{M} \cos \frac{\xi}{2}. \quad (16)$$

Note that symmetry of the reflection under Γ requires

$$R_{\frac{1}{2}\frac{1}{2}}^0(\theta) = R_{\frac{1}{2}\frac{1}{2}}^1(\theta),$$

thus in the first case ($BSSG^+$) the reflections also commute with the operator Γ , while in the other case ($BSSG^-$) they do not. We remark that there are no poles in the physical strip in any of the reflection factors above.

In the case $BSSG^+$, the supersymmetry constraints do not determine the value of γ in contrast to the results of [7, 22]. The reason is that it is the supersymmetry of the reflections on $|B_a\rangle$, $a = 0, 1$ which connects γ with a parameter in the reflection matrix itself. However, we construct these reflections by the bootstrap, which determines them completely, and γ is left as a free parameter. As it was argued above and as will also be seen later γ is connected to the vacuum energy, so it is not a new parameter of the theory (in principle it is expressible in terms of the Lagrangian parameters). The only independent parameters introduced by the boundary are η and ϑ which are present in the bosonic sine-Gordon reflection factors R_{SG} .


5.2 The general spectrum and the associated reflection factors

5.2.1 The Γ symmetric case ($BSSG^+$)

We start with the analysis of the ground state reflection factors

$$R_{\frac{1}{2}\frac{1}{2}}^a(\theta) \times R_{SG}(\theta)$$

where the SUSY component has the form (14). Since the only poles of these reflection factors are due to the sine-Gordon part their explanation has to be similar to that in the bosonic theory. However, we have to supplement the formulae for the bosonic theory with RSOS indices in a consistent way. The sine-Gordon reflection factor has boundary independent poles at $i\frac{n\pi}{2\lambda}$ for $n = 1, 2, \dots$, which can be described by diagram (a). This is identical to the non supersymmetric diagram except that it is decorated with RSOS indices, which are displayed inside circles. Clearly the dashed line denotes the full breather supermultiplet (now consisting of a boson and a fermion).



The boundary dependent poles of R_{SG} are located at

$$-i\theta = \nu_n = \frac{\eta}{\lambda} - \frac{(2n+1)\pi}{2\lambda} \quad ; \quad n = 0, 1, \dots$$

At the position of these poles we associate boundary bound states to the reflection amplitudes $R_{\frac{1}{2}\frac{1}{2}}^a$, $a = 0, 1$

$$|a, 1/2|n\rangle = \frac{1}{g_{|1/2\rangle}^{[1/2]}} K_{a\frac{1}{2}}(i\nu_n) \left| \frac{1}{2} \right\rangle, \quad \text{where } \left| \frac{1}{2} \right\rangle \equiv |B_{\frac{1}{2}}\rangle, \quad (17)$$

where the g -factor is the SUSY part of the boundary coupling, coming from the SUSY component of the reflection factor (for definitions of boundary couplings, see [18]). The two states ($a = 0, 1$) for a given n form a doublet which realizes the structure (11), that is the $K_{\frac{1}{2}a}$ kinks can scatter on it. The action of the boundary supercharge on these states can be calculated using the coproduct rules in [10, 11], taking into account the action of the charges on the boundary ground state:

$$\begin{aligned} \tilde{Q}_+ |0, 1/2|n\rangle &= r \left(\gamma - 2i\sqrt{M} \cos \frac{\nu_n}{2} \right) |1, 1/2|n\rangle \\ \tilde{Q}_+ |1, 1/2|n\rangle &= r^{-1} \left(\gamma + 2i\sqrt{M} \cos \frac{\nu_n}{2} \right) |0, 1/2|n\rangle, \quad r = \frac{g_{|1, 1/2|n\rangle}^{[1/2]}}{g_{|0, 1/2|n\rangle}^{[1/2]}} \end{aligned} \quad (18)$$

The boundary supercharge satisfies

$$\tilde{Q}_+^2 |a, 1/2|n\rangle = 2 \left(\frac{\gamma^2}{2} + M \cos \nu_n + M \right) |a, 1/2|n\rangle$$

which is exactly the relation $\tilde{Q}_+^2 = 2 \left(\tilde{H} + M \tilde{Z} \right)$, since the central charge of this state is $\tilde{Z} = 1$ (we take the ground state $|1/2\rangle$ to have $\tilde{Z} = 0$, while the bulk soliton $K_{a1/2}$ has $Z = 1$), the ground state has energy $\gamma^2/2$ by virtue of the relations (12,13) and $M \cos \nu_n$ is the energy that the excited state has relative to the ground state (10).

The SUSY reflection factors of $K_{\frac{1}{2}a}$ off $|a, 1/2|n\rangle$ can be computed from the bootstrap principle (diagrams (b) and (c)):

$$g_a^{\frac{1}{2}} R_{ab}^{\frac{1}{2}}(\theta) = g_b^{\frac{1}{2}} \left\{ \sum_{x=0,1} S \begin{pmatrix} \frac{1}{2} & x \\ a & \frac{1}{2} \end{pmatrix} (\theta - i\nu_n) S \begin{pmatrix} \frac{1}{2} & b \\ x & \frac{1}{2} \end{pmatrix} (\theta + i\nu_n) R_{\frac{1}{2}\frac{1}{2}}^x(\theta) \right\} \quad (19)$$

where $g_a^{\frac{1}{2}} \equiv g_{|a,1/2|n}^{[1/2]}$

The result turns out to be


$$R_{ab}^{\frac{1}{2}}(\theta) = P(\theta) K(\theta + i\nu_n) K(\theta - i\nu_n) \frac{g_b^{\frac{1}{2}}}{g_a^{\frac{1}{2}}} \left(\delta_{ab} \cos \left(\frac{\nu_n}{2} \right) + \delta_{a,1-b} \sin \left(\frac{\theta}{2i} \right) \right) \quad (20)$$

Note the appearance of the g factors in the result. They are the SUSY parts of the boundary couplings and come in two types: one corresponds to the absorption of the particle while creating a higher excited boundary state, the other describing the emission of the particle and transition to some lower excited boundary state. The ones above are of the absorption type. The residues of the full reflection factor are described by the product of an emission and an absorption type full boundary coupling (for a definition of boundary couplings and their relation to the residue of the reflection factor see [18]). Due to the tensor product structure the full boundary coupling is given by the bosonic part multiplied with the SUSY g -factor, as in the case of bulk scattering [11]. The product of the appropriate emission and absorption SUSY g -factors is constrained to coincide with the value of the SUSY part of the reflection factor at the position of the pole in the bosonic factor. It can be seen in general that this does not give enough constraints to determine their value unambiguously due to the degeneracy introduced by the RSOS indices a, b , and as no physical quantity should explicitly depend on their value (see the example of the relation $\tilde{Q}_+^2 = 2 \left(\tilde{H} + M \tilde{Z} \right)$ discussed above) we do not present any solution for them.

Being constructed by the bootstrap, the reflection factors (20) necessarily satisfy the constraints of boundary factorization and crossing-unitarity; in addition, they commute with $\tilde{Q}_+$ which is guaranteed by the fact that the action of the boundary supercharge is also derived from the bootstrap as in (18). The full reflection factor on the $|a, 1/2|n\rangle$ excited boundary can be obtained by multiplying this result with the appropriate excited bosonic reflection factor:

$$R_{ab}^{\frac{1}{2}}(\theta) \times Q_{|n\rangle}(\eta, \vartheta, \theta) \quad \text{or} \quad R_{ab}^{\frac{1}{2}}(\theta) \times P_{|n\rangle}^{\pm}(\eta, \vartheta, \theta) . \quad (21)$$

Clearly (20) has neither pole nor zero in the physical strip. So the poles of the reflection factors on the first excited wall (21) are exactly the same as in the non supersymmetric theory: that is they are at $i\nu_k$ or at $i\nu_m$.



The decoration of the non supersymmetric diagrams shows, that diagram (e) explains the ν type of poles, while diagram (d) explains the w type, but only for $w_m > \nu_n$. For $w_m < \nu_n$ we have a boundary bound state which we denote by

$$\left| \frac{1}{2}, a, \frac{1}{2} \middle| m, n \right\rangle = \frac{1}{g_{\left| \frac{1}{2}, a, \frac{1}{2} \middle| m, n \right\rangle}} K_{\frac{1}{2}a} (i w_m) |a, 1/2|n\rangle ,$$

so this is also a doublet, but now it is the $K_{a\frac{1}{2}}$ type kinks that are able to reflect on it. It can be checked easily that the relation $\tilde{Q}_+^2 = 2 \left(\tilde{H} + M \tilde{Z} \right)$ holds for these states as well, consistently with the previous interpretation of γ (for these states $\tilde{Z} = 0$).

At this point the question emerges whether the two states $a = 0, 1$ forming the doublet $\left| \frac{1}{2}, a, \frac{1}{2} \middle| m, n \right\rangle$ are physically different or there is a possibility for some identification so that a single state can explain the pole in the reflection matrix (21). This can be decided by examining whether one can describe the residue of the reflection factor with a single intermediate state, which implies a relation between the $R_{00}^{1/2}, R_{11}^{1/2}, R_{10}^{1/2}$ and $R_{01}^{1/2}$ components of the reflection factor at the pole. This relation is violated (for generic values of the parameters) and so one must really introduce the two states above.

Following the same analysis we performed in [21], but now using a decorated version of the Coleman-Thun diagrams it can be seen that the poles in the reflection matrix on the above boundary excited state, which can not be explained by Coleman-Thun diagrams are located at $i\nu_k$. Since the poles appear in association with both reflection factors $R_{\frac{1}{2}\frac{1}{2}}^b(\theta)$, the corresponding boundary states, which are denoted by


$$\left| b, \frac{1}{2}, a, \frac{1}{2} \middle| k, m, n \right\rangle$$

have a fourfold degeneracy.

It is clear that the general boundary bound state has the structure

$$\left| a_k \dots \frac{1}{2}, a_1, \frac{1}{2} \middle| n_k \dots, m_1, n_1 \right\rangle \quad \text{or} \quad \left| \frac{1}{2}, a_k \dots \frac{1}{2}, a_1, \frac{1}{2} \middle| m_k, n_k \dots, m_1, n_1 \right\rangle \quad (22)$$

From this we see that in the supersymmetric case the boundary excited states have a nontrivial degeneracy in contrast to the bosonic theory. The degeneracy is labeled by RSOS sequences starting from $1/2$. In both states in (22) the labels a_i can freely take the values 0 and 1, and, as a result, the degeneracy of the states is 2^k . The associated reflection factors can be computed from successive application of the bootstrap procedure, which is illustrated on the figure



The result depends on the $\tilde{Z}$ charge of the scattering particles. In the $\tilde{Z} = 0$ case the result can be written in the following form

$$R^{\frac{1}{2}}(\theta) \left| b_k \dots \frac{1}{2}, b_1, \frac{1}{2} \middle| n_k \dots, m_1, n_1 \right\rangle = R^{\frac{1}{2}}_{a_1 b_1}(\theta) \prod_{i=1}^{k-1} f^{a_i a_{i+1}}_{b_i b_{i+1}}(w_{m_i}, \nu_{n_{i+1}}, \theta) \quad (23)$$

where $f^{a_1 a_2}_{b_1 b_2}(w_{m_1}, \nu_{n_2}, \theta)$ is the contribution of the dotted square summing over $x_1 = 0, 1$ that is

$$\begin{aligned} f^{a_1 a_2}_{b_1 b_2}(w_{m_1}, \nu_{n_2}, \theta) &= \sum_{x_1=0,1} S \left(\begin{array}{cc} x_1 & \frac{1}{2} \\ \frac{1}{2} & a_1 \end{array} \right) (\theta - i w_{m_1}) S \left(\begin{array}{cc} x_1 & \frac{1}{2} \\ \frac{1}{2} & b_1 \end{array} \right) (\theta + i w_{m_1}) \\ &\times S \left(\begin{array}{cc} \frac{1}{2} & x_1 \\ a_2 & \frac{1}{2} \end{array} \right) (\theta - i \nu_{n_2}) S \left(\begin{array}{cc} \frac{1}{2} & b_2 \\ x_1 & \frac{1}{2} \end{array} \right) (\theta + i \nu_{n_2}) \end{aligned}$$

Collecting the common factors we have

$$\begin{aligned} f^{a_1 a_2}_{b_1 b_2}(w_{m_1}, \nu_{n_2}, \theta) &= K(\theta - i w_{m_1}) K(\theta + i w_{m_1}) K(\theta - i \nu_{n_2}) K(\theta + i \nu_{n_2}) \\ &\times \frac{g^{a_1, \frac{1}{2}}_{b_1, \frac{1}{2}} | n_1 \rangle g^{\frac{1}{2} a_1, \frac{1}{2}} | m_1, n_1 \rangle}{\{a \leftrightarrow b\}} h^{a_1 a_2}_{b_1 b_2}(w_{m_1}, \nu_{n_2}, \theta) \end{aligned}$$

where

$$h_{b_1 b_2}^{a_1 a_2}(w_{m_1}, \nu_{n_2}, \theta) = 2 \sum_{x=0,1} \cos \left(\frac{\theta - i w_{m_1}}{4i} + \frac{\pi}{4}(1 - 2\delta_{x,a_1}) \right) \cos \left(\frac{\theta + i w_{m_1}}{4i} + \frac{\pi}{4}(1 - 2\delta_{x,b_1}) \right) \\ \times \cos \left(\frac{\theta - i \nu_{n_2}}{4i} - \frac{\pi}{2}\delta_{x,a_2} \right) \cos \left(\frac{\theta + i \nu_{n_2}}{4i} - \frac{\pi}{2}(1 - \delta_{x,b_2}) \right)$$

In the $\tilde{Z} = 1$ case, which is indicated with dotted lines on the diagram, the result contains an extra factor

$$R^{x_k}(\theta) \left| \frac{1}{2}, b_k \dots \frac{1}{2}, b_1, \frac{1}{2} \middle| m_k \dots, m_1, n_1 \right\rangle = h_{a_k, b_k}^{x_k} R^{\frac{1}{2}}(\theta) \left| b_k \dots \frac{1}{2}, b_1, \frac{1}{2} \middle| n_k \dots, m_1, n_1 \right\rangle \quad (24)$$

where

$$h_{a_k, b_k}^{x_k} = 2^{-\frac{\theta}{i\pi}} \frac{g \left| \frac{1}{2} a_k \frac{1}{2} \dots \right\rangle}{g \left| \frac{1}{2} b_k \frac{1}{2} \dots \right\rangle} \left(-\sin \left(\frac{\theta}{2i} - \frac{\pi}{2}(\delta_{x_k, a_k} + \delta_{x_k, b_k}) \right) + \cos \left(\frac{w_{m_k}}{2} + \frac{\pi}{2}(\delta_{x_k, a_k} - \delta_{x_k, b_k}) \right) \right)$$

5.2.2 The Γ non symmetric ($BSSG^-$) case

The discussion of the Γ non symmetric ($BSSG^-$) solution runs entirely parallel to the previous Γ symmetric case, the only difference being that in the input of the bootstrap procedure, i.e. in the ground state reflection amplitude

$$R_{\frac{1}{2}\frac{1}{2}}^a(\theta) \times R_{SG}(\theta)$$

the supersymmetry factors, $R_{\frac{1}{2}\frac{1}{2}}^a(\theta)$, $a = 0, 1$, are taken now from (15). These factors depend explicitly on γ , and this dependence pertains in the (SUSY) reflection amplitudes on exited boundaries. Nevertheless, since none of these amplitudes has a pole in the physical strip, following the steps of the previous considerations leads to the same conclusion regarding the indexing and degeneracies of the boundary states. Therefore we concentrate here mainly on the differences between the two solutions.

At the position of the ν_n poles in the ground state reflection amplitude we again associate boundary bound states $|a, 1/2|n\rangle$ to $R_{\frac{1}{2}\frac{1}{2}}^a$ $a = 0, 1$ as in (17) (though of course the present values of boundary couplings may differ from the previous ones). The action of the present boundary supercharge, $\tilde{Q}_-$, on these states is

$$\tilde{Q}_- |0, 1/2|n\rangle = r \left(\gamma + 2\sqrt{M} \sin \frac{\nu_n}{2} \right) |1, 1/2|n\rangle,$$

$$\tilde{Q}_- |1, 1/2|n\rangle = r^{-1} \left(\gamma - 2\sqrt{M} \sin \frac{\nu_n}{2} \right) |0, 1/2|n\rangle.$$

The action of $\tilde{Q}_-^2$ on these states is compatible with the relation $\tilde{Q}_-^2 = 2(\tilde{H} - M\tilde{Z})$, provided we keep the interpretation of $\gamma^2/2$ as the ground state energy.

Using the SUSY factors eqn. (15) in the bootstrap equation (19) for the reflections of the $K_{\frac{1}{2}a}$ kinks on these boundary states gives

$$\begin{aligned} R_{ab}^{\frac{1}{2}}(\theta) &= Z_-(\theta) \frac{g_b^{\frac{1}{2}}}{g_a^{\frac{1}{2}}} \left[\delta_{ab} \left(\cos \frac{\xi}{2} \cos \frac{\nu_n}{2} + (-1)^a i \sinh \frac{\theta}{2} \cosh \frac{\theta}{2} \right) \right. \\ &\quad \left. - i \delta_{a,1-b} \sinh \frac{\theta}{2} \left(\cos \frac{\xi}{2} + (-1)^b \sin \frac{\nu_n}{2} \right) \right], \end{aligned}$$

where ξ is expressed in terms of γ in eqn. (16), and

$$Z_-(\theta) = P(\theta) K(\theta + i\xi) K(\theta - i\xi) K(\theta + i\nu_n) K(\theta - i\nu_n).$$

Although this reflection amplitude has a slightly more complicated form than the one in eqn. (20), it also solves the boundary Yang-Baxter equation and the other constraints by construction. The difference between the two comes from the fact that (20) commutes with $\tilde{Q}_+$, while the present reflection factor is invariant under $\tilde{Q}_-$.

Finally we point out that the expressions (23,24) for the kink reflection amplitudes on the general higher excited boundary states remain valid in the case $BSSG^-$ as well, if the ground state reflection factor (14) is replaced by (15).

6 Breather reflection factors

The breathers have vertex type scattering matrices in contrast to the RSOS type ones of the kinks. These scattering matrices enter into the equations determining the reflection factors of the breathers, nevertheless there is no need for their explicit form as the breather reflection factors on the various boundaries can be obtained from that of the soliton kinks by using the (bulk) fusion and the bootstrap [19]; the procedure is summarized schematically on diagram (f).

If two bulk kinks form a bound state at a rapidity difference $i\rho$ ($0 < \rho < \pi$) the bound state is identified with a supermultiplet (ϕ, ψ) of mass $2M \cos(\rho/2)$. (In case of the k th breather $\rho = \rho_k = \pi - \frac{k\pi}{\lambda}$). The fusing coefficients of these processes are defined via [11]:


$$|K_{ab}(\theta + i\rho/2) K_{bc}(\theta - i\rho/2)\rangle = f_{abc}^{\phi} |\phi(\theta)\rangle + f_{abc}^{\psi} |\psi(\theta)\rangle$$

with the non vanishing coefficients being:

$$f_{0\frac{1}{2}0}^{\phi} = f_{1\frac{1}{2}1}^{\phi} = 2^{(\pi-2\rho)/4\pi} f_{\frac{1}{2}0\frac{1}{2}}^{\phi} = 2^{(\pi-2\rho)/4\pi} f_{\frac{1}{2}1\frac{1}{2}}^{\phi} = \sqrt{K(i\rho) 2^{(\pi-\rho)/2\pi} \cos\left(\frac{\rho-\pi}{4}\right)}$$

and

$$f_{1\frac{1}{2}0}^{\psi} = -f_{0\frac{1}{2}1}^{\psi} = 2^{(\pi-2\rho)/4\pi} i f_{\frac{1}{2}0\frac{1}{2}}^{\psi} = -2^{(\pi-2\rho)/4\pi} i f_{\frac{1}{2}1\frac{1}{2}}^{\psi} = \sqrt{K(i\rho) 2^{(\pi-\rho)/2\pi} \cos\left(\frac{\rho+\pi}{4}\right)}$$



(f) The bootstrap procedure for the breather reflection factors on the boundary ground state $|\frac{1}{2}\rangle$

To describe the ground state reflection amplitudes of the bosonic (ϕ) and fermionic (ψ) components we represent them as

$$\phi(\theta)|\frac{1}{2}\rangle = \frac{1}{2f^{\phi}_{\frac{1}{2}0\frac{1}{2}}} \left(K_{\frac{1}{2}0}(\theta + i\rho/2)K_{0\frac{1}{2}}(\theta - i\rho/2) + K_{\frac{1}{2}1}(\theta + i\rho/2)K_{1\frac{1}{2}}(\theta - i\rho/2) \right) |\frac{1}{2}\rangle,$$

$$\psi(\theta)|\frac{1}{2}\rangle = \frac{1}{2f^{\psi}_{\frac{1}{2}0\frac{1}{2}}} \left(K_{\frac{1}{2}0}(\theta + i\rho/2)K_{0\frac{1}{2}}(\theta - i\rho/2) - K_{\frac{1}{2}1}(\theta + i\rho/2)K_{1\frac{1}{2}}(\theta - i\rho/2) \right) |\frac{1}{2}\rangle.$$

These expressions show that they also provide an ordinary doublet representation of the boundary supercharge $\tilde{Q}_{\pm}$ and that the fermionic parity Γ act on them in the standard way. The actual reflection factors are obtained from the bootstrap equation on diagram (f), where the dashed lines represent either ϕ or ψ . The bosonic and fermionic reflection factors are qualitatively different in the Γ symmetric and Γ non symmetric cases, since the bootstrap equations contain both the $R_{\frac{1}{2}\frac{1}{2}}^0$ and the $R_{\frac{1}{2}\frac{1}{2}}^1$ ground state kink reflection amplitudes, and these are significantly different in the two cases. Writing the breather reflection factors on the ground state boundary as

$$\begin{pmatrix} \phi \\ \psi \end{pmatrix}(\theta)|\frac{1}{2}\rangle = \begin{pmatrix} \mathcal{A}_+ & \mathcal{B} \\ \tilde{\mathcal{B}} & \mathcal{A}_- \end{pmatrix} \begin{pmatrix} \phi \\ \psi \end{pmatrix}(-\theta)|\frac{1}{2}\rangle$$

in the Γ symmetric ($BSSG^+$) case one obtains

$$\mathcal{B} = \tilde{\mathcal{B}} = 0, \quad \mathcal{A}_+ = Z(\theta) \cos\left(\frac{\theta}{2i} - \frac{\pi}{4}\right), \quad \mathcal{A}_- = Z(\theta) \cos\left(\frac{\theta}{2i} + \frac{\pi}{4}\right),$$

$$Z(\theta) = P(\theta + i\rho/2)P(\theta - i\rho/2)\sqrt{2}K(2\theta)2^{-\theta/i\pi} ,$$

while in the Γ non symmetric ($BSSG^-$) case we get

$$\mathcal{B} = -\tilde{\mathcal{B}} = -\tilde{Z}(\theta)\frac{\gamma}{2\sqrt{M}}\sqrt{\cos(\rho/2)}\sinh(\theta),$$

$$\begin{aligned} \mathcal{A}_{\pm} = \tilde{Z}(\theta) & \left(\cosh\left(\frac{\theta}{2}\right) \left(\frac{\gamma^2}{4M} - \left[\sin^2\left(\frac{\rho}{4}\right) + \sinh^2\left(\frac{\theta}{2}\right) \right] \right) \right. \\ & \left. \mp i \sinh\left(\frac{\theta}{2}\right) \left(\frac{\gamma^2}{4M} + \left[\sin^2\left(\frac{\rho}{4}\right) + \sinh^2\left(\frac{\theta}{2}\right) \right] \right) \right) , \end{aligned}$$

with

$$\tilde{Z}(\theta) = K(2\theta)2^{-\theta/i\pi}F(\theta - i\rho/2)F(\theta + i\rho/2), \quad F(\theta) = P(\theta)K(\theta + i\xi)K(\theta - i\xi) .$$

The two cases are indeed qualitatively different: in the $BSSG^+$ solution the conservation of fermionic parity forbids the $\phi \rightarrow \psi$ reflection on the ground state boundary, while in the $BSSG^-$ solution this reflection is possible. The form of the $BSSG^+$ solution is the same as the one obtained in [24] by imposing fermion number conservation. The structural form of the reflection factors in the case $BSSG^-$ are identical to the ones obtained in [23] from the eight vertex free fermion model with boundary. In [4] it was proposed that γ , which appears explicitly in the reflection matrix, could be fixed in terms of η and ϑ by the boundary bootstrap. Here we see that this is not the case, as the bootstrap gives no constraint for γ , due to degeneracies appearing in the boundary excited states. We recall however that γ is not a free parameter, as it is determined by the ground state boundary energy (or equivalently, by the boundary supercharge), thus it can be expressed in terms of the Lagrangian parameters in principle.

A more explicit description of the reflection factor of the first breather on the ground state boundary in the $BSSG^-$ case was given in boundary sinh-Gordon model studied in [6], but the precise connection between the parameters used in that paper and the present one is yet to be determined.

In the bosonic theory the ground state reflection amplitude of the k th breather has poles at

$$-i\theta = \frac{\eta}{\lambda} - \frac{\pi}{2} + (k - 2l - 1)\frac{\pi}{2\lambda} \quad , \quad l = 0, \dots, \left\lfloor \frac{k-1}{2} \right\rfloor .$$

In the supersymmetric theory, these poles signal the presence of the excited boundary states

$$\left| \frac{1}{2}, a, \frac{1}{2} \middle| l, k-l \right\rangle$$

as intermediate states in the breather reflection process. Since there are two intermediate states ($a = 0, 1$), the determinant of the 2×2 reflection matrix should not vanish at the position of these poles (as the residue of the reflection matrix at the pole is proportional

to the projector on the subspace of on-shell intermediate states). It is straightforward to verify that this is indeed the case for both the $BSSG^+$ and the $BSSG^-$ solutions.

Using the bootstrap procedure it is also possible to obtain the breather reflection factors on excited boundaries. If the RSOS sequence characterizing the boundary state ends with a label a ($a = 0$ or $a = 1$), then e.g. the bosonic breather can be represented by

$$\left| \phi(\theta)a, \frac{1}{2}, \dots \left| n_1, \dots \right. \right\rangle = \frac{1}{f^\phi_{a\frac{1}{2}a}} K_{a\frac{1}{2}}(\theta + i\rho/2) K_{\frac{1}{2}a}(\theta - i\rho/2) \left| a, \frac{1}{2}, \dots \left| n_1, \dots \right. \right\rangle$$

in the bootstrap procedure. To emphasize that even in the $BSSG^+$ case there are $\phi \rightarrow \psi$ type reflections on excited boundaries we give here the reflection matrix of breathers on the $|a1/2|n\rangle$ states. In the basis of $|\phi(\theta)0, 1/2|n\rangle$, $|\phi(\theta)1, 1/2|n\rangle$, $|\psi(\theta)0, 1/2|n\rangle$, $|\psi(\theta)1, 1/2|n\rangle$ it can be written as

$$\hat{Z}(\theta) \begin{pmatrix} \mathcal{C}_+ & 0 & 0 & -\mathcal{D}/r \\ 0 & \mathcal{C}_+ & r\mathcal{D} & 0 \\ 0 & \mathcal{D}/r & \mathcal{C}_- & 0 \\ -r\mathcal{D} & 0 & 0 & \mathcal{C}_- \end{pmatrix} \quad (25)$$

where $\mathcal{D} = \frac{1}{\sqrt{2}} \sqrt{\cos \frac{\rho}{2}} \cos \frac{\nu_n}{2} \sin \frac{\theta}{i}$ and

$$\mathcal{C}_\pm = \cos^2 \frac{\nu_n}{2} \cos \left(\frac{\theta}{2i} \mp \frac{\pi}{4} \right) + \frac{1}{2} \left(\cos \frac{\rho}{2} - \cos \frac{\theta}{i} \right) \cos \left(\frac{\theta}{2i} \pm \frac{\pi}{4} \right).$$

It is easy to show that in spite of the non trivial boson fermion reflection the operator Γ commutes with this reflection matrix.

A nontrivial check on the consistency of the bootstrap solution can be obtained by considering the pole structure of the full reflection amplitude containing the SUSY factor (25). The bosonic reflection factor of $\mathbf{B}^k$ on the bosonic boundary excited state $|n\rangle$ has a pole at $-i\theta = \frac{\pi}{2} - \frac{\eta}{\lambda} + \frac{\pi}{2\lambda} (k + 2n + 1)$ [21]. In the supersymmetric case, this means that a boundary excited state of the form

$$\left| a, \frac{1}{2} \left| n + k \right. \right\rangle \quad (26)$$

enters as an on-shell intermediate state in the scattering of $\mathbf{B}^k$ on $|a, 1/2|n\rangle$. However, due to the doublet (boson/fermion) structure of the breather naively one would expect 4 states to explain the residue of the 4×4 reflection factor. In the conjectured spectrum, on the other hand, the only possible process goes via (26) and it allows for only two intermediate states ($a = 0, 1$). Therefore one expects that the determinant of the matrix (25) should have a double zero there. It can be verified by direct calculation that this double zero is indeed there without imposing any restriction on the parameters.

In the reflection of the k th breather on the n th excited boundary, there is another family of poles at $-i\theta = \frac{\eta}{\lambda} - \frac{\pi}{2\lambda} (k - 2l + 1)$, $l = 0, \dots, n - 1$ [21] that in the supersymmetric case should correspond to intermediate states of the form

$$\left| b, \frac{1}{2}, a, \frac{1}{2} \left| l, k - l, n \right. \right\rangle.$$

At these poles, the number of intermediate states is 4 ($a, b = 0, 1$) and so we expect that the determinant of the SUSY factor does not vanish, which indeed turns out to be the case.

7 Conclusions

To start we summarize the results of this paper. We considered the boundary scattering amplitudes in boundary supersymmetric sine-Gordon theory (*BSSG*). Imposing the constraint of supersymmetry on solutions of the boundary Yang-Baxter equation, we found two consistent sets of amplitudes that describe the reflection of solitons off the boundary in its ground state. Then we considered the two bootstrap systems built from these fundamental amplitudes and conjectured the closure of this bootstrap, i.e. the set of boundary states and the reflection factors on them. We also derived a relation between the boundary supercharge and the Hamiltonian and checked that this relation holds for the bootstrap solutions.

Although the reflection amplitudes are different, the spectrum of states is the same in the two bootstrap solutions. This common spectrum is characterized partly by a sequence of integers, just like in the case of the ordinary sine-Gordon model [21], but also by an RSOS sequence of length $k+1$ (if the length of the integer sequence is k) starting from $1/2$. The energy of the state depends only on the integer labels, the different RSOS sequences correspond to degenerate states. It is interesting to note that the non supersymmetric boundary spectrum allows for a tensor product type supersymmetrization, and no further constraints are obtained in accord with the bulk case [11].

In the case of the $BSSG^+$ theory, the reflection amplitudes depend on two parameters η and ϑ , that are inherited from the bosonic reflection factors and were originally introduced in [18]. In the bosonic case it is known how these parameters are related to the parameters of the boundary Lagrangian in the perturbed CFT formalism [25]. Besides that, the SUSY algebra introduces a further parameter γ , which is related to the energy of the boundary ground state and so must be a function of the parameters of the *BSSG* Lagrangian. In the $BSSG^-$ theory the difference is that γ appears also in the expression for the reflection factors themselves. In the bosonic case the expression for the boundary energy in terms of Lagrangian parameters is also known [25]. The existence of two different families of solutions and the number of parameters are in accordance with the expectations that they describe the scattering in the Lagrangian theories corresponding to the boundary interaction (6) [4].

It is a very interesting and important issue to connect the bootstrap parameters η , θ and the vacuum energy parameter γ to the parameters of the Lagrangian description for the supersymmetric case as well. In the case of the non supersymmetric boundary sine-Gordon theory this was achieved by considering it as a combined bulk and boundary perturbation of a $c = 1$ free massless boson with Neumann boundary condition. However, even the interpretation of the bulk *SSG* theory as a perturbed CFT is nontrivial, and we are investigating this problem. We are also working on getting more evidence to link the bulk S matrix and the reflection factors to the Lagrangian theory. Work is in progress in

these directions and we hope to report on the results in the very near future.

Acknowledgments

The authors would like to thank G.M.T. Watts for very useful discussions and R.I. Nepomechie for comments on the manuscript. G.T. thanks the Hungarian Ministry of Education for a Magyary Postdoctoral Fellowship, while B.Z. acknowledges partial support from a Bolyai János Research Fellowship. This research was supported in part by the Hungarian Ministry of Education under FKFP 0043/2001 and the Hungarian National Science Fund (OTKA) grants T037674/02 and T34299/01.

References

- [1] S. Ferrara, L. Girardello and S. Sciuto: **An infinite set of conservation laws of the supersymmetric sine-Gordon theory**, *Phys. Lett.* **B76** (1978) 303.
- [2] C. Ahn: **Complete S-matrices of supersymmetric sine-Gordon theory and perturbed superconformal minimal model**, *Nucl. Phys.* **B354** (1991) 57-84.
- [3] T. Inami, S. Odake and Y.-Z. Zhang: **Supersymmetric extension of the sine-Gordon theory with integrable boundary interactions**, *Phys. Lett.* **B359** (1995) 118-124, hep-th/9506157.
- [4] R.I. Nepomechie: **The boundary supersymmetric sine-Gordon model revisited**, *Phys. Lett.* **B509** (2001) 183-188, hep-th/0103029.
- [5] C. Ahn and W.M. Koo: **Exact boundary S matrices of the supersymmetric sine-Gordon theory on a half line**, *J. Phys.* **A29** (1996) 5845-5854, hep-th/9509056.
- [6] C. Ahn and R.I. Nepomechie: **Exact solution of the supersymmetric sinh-Gordon model with boundary**, *Nucl. Phys.* **B586** (2000) 611-640, hep-th/0005170.
- [7] R.I. Nepomechie: **Supersymmetry in the boundary tricritical Ising field theory**, preprint UMTG-234, hep-th/0203123
- [8] A.B. Zamolodchikov: **Fractional-spin integrals of motion in perturbed conformal field theory**, in *Fields, Strings and Quantum Gravity*, eds. H. Guo, Z. Qiu and H. Tye, (Gordon and Breach, 1989).
- [9] R. Shankar and E. Witten: **The S matrix of the supersymmetric nonlinear sigma model**, *Phys. Rev.* **D17** (1978) 2134
- [10] K. Schoutens: **Supersymmetry and factorizing scattering**, *Nucl. Phys.* **B344** (1990) 665-695.

- [11] T.J. Hollowood and E. Mavrikis: The $N=1$ supersymmetric bootstrap and Lie algebras, *Nucl. Phys.* **B484** (1997) 631-652, hep-th/9606116.
- [12] G. Takács: A new RSOS restriction of the Zhiber-Mikhailov-Shabat model and $\Phi(1,5)$ perturbations of nonunitary minimal models, *Nucl. Phys.* **B489** (1997) 532-556, hep-th/9604098.
- [13] C. Ahn and R.I. Nepomechie: The scaling supersymmetric Yang-Lee model with boundary, *Nucl. Phys.* **B594** (2001) 660-684, hep-th/0009250.
- [14] E. Witten and D.I. Olive: Supersymmetry algebras that include topological charges, *Phys. Lett.* **B78** (1978) 97.
- [15] Z. Bajnok, L. Palla, G. Takács and F. Wágner: The k -folded sine-Gordon model in finite volume, *Nucl. Phys.* **B587** (2000) 585-618, hep-th/0004181.
- [16] N. Graham and R.L. Jaffe: Energy, central charge, and the BPS bound for $(1+1)$ -dimensional supersymmetric solitons, *Nucl. Phys.* **B544** (1999) 432-447, hep-th/9808140.
M.A. Shifman, A.I. Vainshtein and M.B. Voloshin: Anomaly and quantum corrections to solitons in two-dimensional theories with minimal supersymmetry, *Phys. Rev.* **D59** (1999) 045016, hep-th/9810068.
A. Litvintsev and P. van Nieuwenhuizen: Once more on the BPS bound for the SUSY kink, preprint YITP-00-18, hep-th/0010051.
A. Rebhan, P. van Nieuwenhuizen and R. Wimmer: The anomaly in the central charge of the supersymmetric kink from dimensional regularization and reduction, preprint TUW-02-16, YITP-02-35, hep-th/0207051.
- [17] R. Chatterjee and A. B. Zamolodchikov: Local magnetization in critical Ising model with boundary magnetic field, preprint RU-93-54, hep-th/9311165.
- [18] S. Ghoshal and A.B. Zamolodchikov: Boundary S matrix and boundary state in two-dimensional integrable quantum field theory, *Int. J. Mod. Phys.* **A9** (1994) 3841-3886, Erratum-ibid. **A9** (1994) 4353, hep-th/9306002.
- [19] S. Ghoshal: Bound state boundary S matrix of the sine-Gordon model, *Int. J. Mod. Phys.* **A9** (1994) 4801, hep-th/9310188.
A. Fring, R. Köberle: Factorized scattering in the presence of reflecting boundaries, *Nucl. Phys.* **B421** (1994) 159, hep-th/9304141.
- [20] P. Mattsson and P. Dorey: Boundary spectrum in the sine-Gordon model with Dirichlet boundary conditions, *J. Phys.* **A33** (2000) 9065-9094, hep-th/0008071.
P. Mattsson Integrable Quantum Field Theories, in the Bulk and with a Boundary, Ph.D. thesis, hep-th/0111261.

- [21] Z. Bajnok, L. Palla, G. Takács and G.Z. Tóth: The spectrum of boundary states in sine-Gordon model with integrable boundary conditions, *Nucl. Phys.* **B622** (2002) 548-564, hep-th/0106070.
- [22] L. Chim: Boundary S matrix for the tricritical Ising model, *Int. J. Mod. Phys.* **A11** (1996) 4491-4512, hep-th/9510008.
- [23] C. Ahn and W.M. Koo: Supersymmetric Sine-Gordon Model and the Eight-Vertex Free Fermion Model with Boundary, *Nucl. Phys.* **B482** (1996) 675, hep-th/9606003.
- [24] M. Moriconi and K. Schoutens: Reflection matrices for integrable N=1 supersymmetric theories, *Nucl. Phys.* **B487** (1997) 756-778, hep-th/9605219.
- [25] Al.B. Zamolodchikov, unpublished.
 Z. Bajnok, L. Palla and G. Takács: Finite size effects in boundary sine-Gordon theory, *Nucl. Phys.* **B622** (2002) 565-592, hep-th/0108157.